

CLAIMS

What is claimed is:

- Sub. 92
1. An antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor. 5
 2. An antibody or antigen-binding fragment according to Claim 1 wherein said antibody or antigen-binding fragment thereof inhibits one or more functions associated with binding of the ligand to the receptor.
 3. An antibody or antigen-binding fragment thereof according to Claim 1 wherein the mammalian CC-chemokine receptor 2 is a human CC-chemokine receptor 2. 10
 4. An antibody or antigen-binding fragment thereof according to Claim 1 wherein the ligand is a chemokine.
 5. An antibody or antigen-binding fragment thereof according to Claim 4 wherein the chemokine is selected from the group consisting of MCP-1, MCP-2, MCP-3, MCP-4 and combinations thereof. 15
 6. An antibody or antigen-binding fragment thereof according to Claim 1, wherein said antibody or fragment is a monoclonal antibody or fragment thereof.
 7. An antibody or antigen-binding fragment thereof according to Claim 1, wherein said antibody or fragment is a human antibody or fragment thereof.

8. An antibody or antigen-binding fragment thereof according to Claim 1, wherein said antigen-binding fragment is selected from the group consisting of an Fv fragment, an Fab fragment, an Fab' fragment and an F(ab')₂ fragment.

9. An antibody or antigen-binding fragment thereof according to Claim 1, wherein said antibody or fragment is a humanized antibody or fragment thereof.

10. An antibody or antigen-binding fragment thereof according to Claim 1, wherein said antibody or fragment is a recombinant antibody or fragment thereof.

11. A test kit for use in detecting the presence of a mammalian CC-chemokine receptor 2 in a biological sample comprising

- a) at least one antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor; and
- b) one or more ancillary reagents suitable for detecting the presence of a complex between said antibody or antigen-binding fragment thereof and said mammalian CC-chemokine receptor 2 or a portion thereof.

12. A method of inhibiting the interaction of a cell bearing mammalian CC-chemokine receptor 2 with a ligand thereof, comprising contacting said cell with an effective amount of an antibody or antigen-binding fragment thereof which binds to mammalian CC-chemokine receptor 2 and inhibits binding of said ligand to the receptor.

13. A method according to Claim 12 wherein the cell is selected from the group consisting of lymphocytes, monocytes, granulocytes, T cells, basophils, and cells comprising a recombinant nucleic acid encoding CCR2 or a portion thereof.

Sub. 92
09898513 .070301

14. A method according to Claim 13 wherein the T cells are selected from the group consisting of CD8+ cells, CD25+ cells, CD4+ cells and CD45RO+ cells.
15. A method according to Claim 12 wherein the ligand is a chemokine.
16. A method according to Claim 15 wherein the chemokine is selected from the group consisting of MCP-1, MCP-2, MCP-3, MCP-4 and combinations thereof.
17. A method of inhibiting HIV infection of a cell, comprising contacting a cell with an effective amount of a composition comprising an antibody or antigen-binding fragment thereof which binds to mammalian CC-chemokine receptor 2 and inhibits HIV entry into said cell.
18. A method according to Claim 17 wherein the cell is selected from the group consisting of lymphocytes, monocytes, macrophages, granulocytes, T cells, and cells comprising a recombinant nucleic acid encoding CCR2 or a portion thereof.
19. A method according to Claim 18 wherein the T cells are selected from the group consisting of CD8+ cells, CD25+ cells, CD4+ cells and CD45RO+ cells.
20. A method of treating HIV in a patient comprising administering to the patient a composition comprising an effective amount of an antibody or antigen-binding fragment thereof which binds to mammalian CC-chemokine receptor 2 and inhibits HIV entry into said cell.
21. A method of detecting expression of mammalian CC-chemokine receptor 2 or portion thereof by a cell, comprising:

- 5 a) contacting a composition comprising a cell to be tested with an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor; and
- b) detecting binding of said antibody or antigen-binding fragment thereof, wherein the binding of said antibody or antigen-binding fragment thereof indicates the presence of said receptor or portion of said receptor on said cell.

22. The method of Claim 21 wherein the composition is a sample comprising human cells.

10 23. A method of detecting a mammalian CC-chemokine receptor 2 or portion of said receptor, comprising:

- 15 a) contacting a sample to be tested with an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor under conditions appropriate for binding of said antibody or fragment thereof thereto; and
- b) detecting or measuring binding of said antibody or antigen-binding fragment thereof,

20 wherein the binding of said antibody or antigen-binding fragment thereof to material in said sample is indicative of the presence of a mammalian CC-chemokine receptor 2 or portion of said receptor in said sample.

24. A method according to Claim 23, wherein the sample is a cellular fraction which, in normal individuals, comprises a mammalian CC-chemokine receptor 2 or portion of said receptor.

09898513.070301

25. ✓ A method of inhibiting a function associated with binding of a chemokine to a mammalian CC-chemokine receptor 2 or a functional portion of said receptor, comprising contacting a composition comprising the receptor or portion with an effective amount of an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody inhibits binding of said chemokine to mammalian CC-chemokine receptor 2 and inhibits one or more functions associated with binding of the ligand to the receptor.
26. A method according to Claim 25 wherein the chemokine is selected from the group consisting of MCP-1, MCP-2, MCP-3, MCP-4 and combinations thereof.
- 10 27. ✓ A method of detecting or identifying an agent which binds a mammalian CC-chemokine receptor 2 or ligand-binding variant thereof, comprising combining
- a) an agent to be tested;
 - b) an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor; and
 - 15 c) a composition comprising a mammalian CC-chemokine receptor 2 or a ligand-binding variant thereof,
- under conditions suitable for binding of said antibody or antigen-binding fragment to said mammalian CC-chemokine receptor 2 or ligand-binding variant thereof, and detecting or measuring binding of said antibody or antigen-binding fragment to said mammalian CC-chemokine receptor 2 or ligand-binding variant thereof.
- 20
28. A method according to Claim 27 wherein the formation of a complex between said antibody or antigen-binding fragment and said mammalian CC-chemokine receptor 2 or ligand-binding variant is monitored, and wherein a decrease in the
- 25

09898513-070301

amount of complex formed relative to a suitable control is indicative that the agent binds said receptor or ligand-binding variant thereof.

29. A method according to Claim 27 wherein the composition comprising a mammalian CC-chemokine receptor 2 or a ligand-binding variant thereof is a cell bearing recombinant CC-chemokine receptor 2 or ligand-binding variant thereof.
30. A method according to Claim 29, wherein the composition comprising a mammalian CC-chemokine receptor 2 or a ligand-binding variant thereof is a membrane fraction of said cell bearing recombinant CC-chemokine receptor 2 or ligand-binding variant thereof.
31. A method according to Claim 27 wherein the antibody or antigen-binding fragment thereof is labeled with a label selected from the group consisting of a radioisotope, spin label, antigen label, enzyme label, fluorescent group and chemiluminescent group.
32. A method according to Claim 27 wherein the agent is an antibody having specificity for a mammalian CC-chemokine receptor 2 or antigen-binding fragment thereof.
33. ✓ A method of inhibiting HIV infection in a patient, comprising administering to the patient a composition comprising an effective amount of an antibody or antigen-binding fragment thereof which binds to mammalian CC-chemokine receptor 2 and inhibits binding of HIV to the receptor.
34. ✓ A method of inhibiting leukocyte trafficking in a patient, comprising administering to the patient a composition comprising an effective amount of an

109898513.070301

-58-

antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2 and inhibits binding of a ligand to the receptor.

35. A method according to Claim 34 wherein the ligand is a chemokine.

5 36. A method according to Claim 35 wherein the chemokine is selected from the group consisting of MCP-1, MCP-2, MCP-3, MCP-4 and combinations of the foregoing.

10 37. A composition comprising an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor, and an optional physiologically acceptable vehicle.

38. An antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor with an IC_{50} of less than about $1.0 \mu\text{g/ml}$.

15 39. An antibody or antigen-binding fragment thereof according to Claim 38 wherein the IC_{50} is less than about $0.05 \mu\text{g/ml}$.

20 40. An antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor, and wherein the antibody or antigen-binding fragment binds the receptor with an affinity of at least about $0.1 \times 10^{-9} \text{ M}$.

05898513.070301

Sub: 92

-59-

41. An antibody or antigen-binding fragment thereof according to Claim 40, wherein the affinity is at least about 1×10^{-9} M.
42. An antibody or antigen-binding fragment thereof according to Claim 40, wherein the affinity is at least about 3×10^{-9} M.
- 5 43. A method of treating a CC-chemokine receptor 2-mediated disorder in a patient, comprising administering to the patient an effective amount of an antibody or antigen-binding fragment thereof which binds to a mammalian CC-chemokine receptor 2, wherein said antibody or antigen-binding fragment thereof inhibits binding of a ligand to the receptor.

FOE020" ET586860

Sub. 2

add a 3